

Pl 1171382

REC'D 24 MAY 2004

WIPO

PCT

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office

May 19, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/474,094

FILING DATE: May 29, 2003

RELATED PCT APPLICATION NUMBER: PCT/US04/07931

By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS



L. Edelen

L. EDELEN
Certifying Officer

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

BEST AVAILABLE COPY

05/29/03

JC860 U.S. PTO

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Approved for use through 10/31/2002. OMB 0851-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No.

EV333282775US

U.S. PTO
60/474094

| INVENTOR(S) | | | | | |
|---|--|---------------------------|--------------|---|-------|
| Given Name (first and middle (if any)) | | Family Name or Surname | | Residence (City and either State or Foreign Country) | |
| Shawn | | DeFrees | | 126 Filly Drive, North Wales, PA 19454 | |
| <input type="checkbox"/> Additional inventors are being named on the _____ separately numbered sheets attached hereto | | | | | |
| TITLE OF THE INVENTION (500 characters max) | | | | | |
| COMPOSITIONS AND METHODS FOR THE PREPARATION AND CONJUGATION OF BIANTENARY POLYMERS INCLUDING POLYETHYLENE GLYCOL | | | | | |
| Direct all correspondence to: | | | | | |
| CORRESPONDENCE ADDRESS | | | | | |
| <input type="checkbox"/> Customer Number | | Type Customer Number here | | Place Customer Number Bar Code Label here | |
| OR | | | | | |
| <input checked="" type="checkbox"/> Firm or Individual Name | | Neose Technologies, Inc. | | | |
| Address | | Rachel Rondinelli | | | |
| Address | | 102 Witmer Road | | | |
| City | | Horsham | | State | PA |
| Country | | Telephone | 215-315-9103 | ZIP | 19044 |
| | | Fax | 215-315-9400 | | |
| ENCLOSED APPLICATION PARTS (check all that apply) | | | | | |
| <input checked="" type="checkbox"/> Specification | | Number of Pages | | 15 | |
| <input type="checkbox"/> Drawing(s) | | Number of Sheets | | <input type="checkbox"/> CD(s), Number | |
| <input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76 | | | | <input checked="" type="checkbox"/> Other (specify) | |
| | | | | return postcard | |
| METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT | | | | | |
| <input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. | | | | FILING FEE AMOUNT (\$) | |
| <input type="checkbox"/> A check or money order is enclosed to cover the filing fees | | | | | |
| <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: | | 502311 | | \$80.00 | |
| <input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached. | | | | | |
| The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government. | | | | | |
| <input checked="" type="checkbox"/> No. | | | | | |
| <input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: | | | | | |

Respectfully submitted,

SIGNATURE



Date 05/29/2003

TYPED or PRINTED NAME Rachel H. Rondinelli

REGISTRATION NO.

45,052

TELEPHONE (215) 325-9103

(if appropriate)

Docket Number:

NEO00262C

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is used by the public to file (and by the PTO to process) a provisional application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, Washington, D.C. 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Box Provisional Application, Assistant Commissioner for Patents, Washington, D.C. 20231.

NEO00262C PR

COMPOSITIONS AND METHODS FOR THE PREPARATION AND CONJUGATION OF BIANTENNARY POLYMERS INCLUDING POLYETHYLENE GLYCOL

This application describes compositions and methods for the preparation and use of biantennary polymers as well as the preparation of mono-dispersed polyethylene glycols (PEGs) and their activated forms. The invention is now described with reference to the following non-limiting Examples and schemes. These Examples and schemes are provided for the purpose of illustration only and the invention should in no way be construed as being limited to these Examples or schemes, but rather should be construed to encompass any and all variations which become evident as a result of the teaching provided herein.

I. Compositions and methods for the preparation and conjugation of biantennary polymers

This application describes compositions and methods for the preparation and use of biantennary polymers. The biantennary structure is generated by conjugating the polymer of interest to a small trifunctional ligand in either a step-wise manner or in one pot. Examples of trifunctional ligands that can be used in this invention are described in Scheme 1. The more exemplary ligands include epichlorohydrin, 1,3-dibromo-2-propanol, ornithine, glutamate and aspartate. The chemistries of conjugation are well known in the art and include activating such groups as hydroxyl, amine, carboxylate via chemical means to create leaving groups for the subsequent reaction with the polymer. Alternatively, the polymer can be activated and conjugated to the trifunctional ligand.

Exemplary polymers that can be conjugated to create a biantennary structure include PEG (polyethyleneglycol), mPEG (methoxypolyethyleneglycol), mPPG (methoxy-polypropyleneglycol), polysialic acid, polyglutamate, polyaspartate, polylactate and the like. These polymers can be prepared as heterodispersed (polydispersed) or monodispersed forms and used in the conjugation procedures, Scheme 2. The heterodispersed mPEG's are prepared by a variety of reported methods with degrees of polymerization ranging from 1 to 20,000 ethylene oxide units. Typically, the mPEGs are separated by size exclusion methodologies and fractionated into ranges of molecular weights. Typically, these ranges are from hundreds to thousands of mass units depending on the size of the PEG. Alternatively, the mPEG is monodispersed, a single molecular weight form, and is prepared by direct chemical synthesis or by separation of a single molecular weight from the polydispersed PEGs.

The activation of the biantennary polymers for conjugation to various ligands can be performed using methods that are standard in the art. For example, a hydroxyl group on the biantennary polymer is

NEO00262C PR


activated with activated forms of carbonate such as the bis-NHS, bis-HOBt or bis-HOAt esters. After activation, the biantennary polymers is conjugated to any suitable ligand such as a protein, nucleotide sugar, peptide, lipid, sugar, DNA, RNA or the like. Exemplary examples are shown in **Scheme 3**.

An example of how to prepare a biantennary PEG is shown in **Scheme 3a**. The method begins with epichlorohydrin and reacts with mPEG under basic reaction conditions. After isolation of the product of the reaction, the biantennary mPEG is activated with the 1,1-bis-HOBt-carbonate to form the biantennary-mPEG-carbonate ester. Reaction of this ester with a nucleotide-sugar then provides a reactant that can be transferred to a protein or glycoprotein by using the appropriate glycosyltransferase.

Additional exemplary embodiments of bi-antennary structures are also described in **Scheme 4** and **Scheme 5**.

Scheme 1. Starting Materials.

a.




X and Y (independently selected) from OR_1 , NR_1R_2 , SR_1 , alkyl-X, aryl-X, alkylaryl-X, branched alkyl-X, $COOR_1$, $CONR_1R_2$.

Z is selected from OR_1 , NR_1R_2 , SR_1 , alkyl $COOR_1$, aryl $COOR_1$, alkylaryl-X, $COOR_1$, $CONR_1R_2$.


R_1 and R_2 (independently selected) from H; alkyl; aryl; branched alkyl; R_1 - R_2 as cyclic ring-aromatic, heteroaromatic or alkyl; activating group; halide; leaving group.

Y-Z (independently selected) from a ring such as epoxide, aziridine, cyclic-sulfonate.

Exemplary examples:



b.




X and Y (independently selected) from OR_1 , NR_1R_2 , SR_1 , alkyl-X, aryl-X, alkylaryl-X, branched alkyl-X, $COOR_1$, $CONR_1R_2$, except when X is NH_2 and Y is $(CH_2)_3NH_2$.

Z is selected from OR_1 , NR_1R_2 , SR_1 , alkyl $COOR_1$, aryl $COOR_1$, alkylaryl-X, $COOR_1$, $CONR_1R_2$.

R_1 and R_2 (independently selected) from H; alkyl; aryl; branched alkyl; R_1 - R_2 as cyclic ring-aromatic, heteroaromatic or alkyl; activating group; halide; leaving group.

Y-Z (independently selected) from a ring such as epoxide, aziridine, cyclic-sulfonate.


Exemplary examples:



NEO00262C PR

Scheme 2. Branched PEG's.

a.






X and Y (independently selected) from OR₁, NR₁R₂, SR₁, alkyl-X, aryl-X, alkylaryl-X, branched alkyl-X, COOR₁, CONR₁R₂.

Z is selected from OR₁, NR₁R₂, SR₁, alkylCOOR₁, arylCOOR₁, alkylaryl-X, COOR₁, CONR₁R₂.

R₁ and R₂ (independently selected) from H; alkyl; aryl; branched alkyl; R₁-R₂ as cyclic ring-aromatic, heteroaromatic or alkyl; activating group; mPEG, PEG, mPPG, polysialic acid, polyglutamate, polyaspartate, polylysine, polyethylenimine, polylactide, polyglyceride, functionalized PEG, polymer.

PEG is polyethyleneglycol; mPEG is methoxypolyethyleneglycol; mPPG is methoxypolypropyleneglycol;

Exemplary examples:




a, b, n and m (independently selected) from 1 to 20,000.

NEO00262C PR

Scheme 3. Activated and Coupled Biantennary Polymers.

a. Exemplary Examples.




b. Exemplary Examples.




Scheme 3. Activated and Coupled Biantennary Polymers.

c. Exemplary Example.



Scheme 4. Additional Bi-antennary structures.




wherein $X = \text{OH}, \text{H}$,
an activating
group, or a
nucleotide sugar;

and $n = 1-10$;

and polymer = PEG, PPG,
mPEG, mPPG, alkyl-PEG,
polyglutamate, polysialic
acid, and the like.


Specific examples include, but are not limited to:



wherein $0 = 1-10,000$;

$m = 1-10,000$;

and $Z = \text{OH}, \text{H}$, an activating
group such as HOBt, HOAT,
N-hydroxysuccinimide ester,
p-nitrophenol ester, and the like.




wherein $m, 0 =$ independent
selected from 1-29,

$a, b =$ independently
selected from 1-24.

and $Z = \text{OH}, \text{H}$, an activating group
such as HOBt, HOAT, N-hydroxy-
succinimide ester, p-nitrophenol
ester, and the like.

Scheme 5. Additional Bi-antennary Polymers:




wherein X, Y = independently selected from O, N, S ;

a = 1-10 ;

Z = OH, H, an activating group, a nucleotide sugar

and polymer = PEG, PPG, mPEG, mPPG, alkyl-PEG, polyglutamic acid, polysialic acid, and the like,


Specific examples include but are not limited to:



wherein m, o = independently selected from 1-20,000 ;

a = 1-10 ; and

Z = OH, H, an activating group such as HOBT, HOA, N-hydroxysuccinimide ester, p-nitrophenol ester, and the like,




wherein m, o = independently selected from 1-20,000 ;

a = 1-10 ; and


Z = OH, H, an activating group such as HOBT, HOAT, N-hydroxysuccinimide ester, p-nitrophenol ester, and the like

Scheme 5 continued. Additional Bi-antennary Polymers



wherein $m, o =$ independently selected from 1-20,000; $a = 1-10$; $b = 1-24$; and $Z = \text{OH}, \text{H}$, an activating group such as HOBT, HOAT, N-hydroxysuccin ester, p-nitrophenol ester, and the like.


and






wherein $m, o =$ independently selected from 1-20,000; $a = 1-10$; and $Z = \text{OH}, \text{H}$, an activating group such as HOBT, HOAT, N-hydroxysuccin ester, p-nitrophenol ester, and the like.

II. Preparation of mono-dispersed polyethylene glycols (PEGs) and their activated forms

This application also describes preparation of mono-dispersed polyethylene glycols (PEGs) and their activated forms. Mono-dispersed or singular molecular weight PEGs can be prepared as shown below. By adjusting the size of the fragments generated, any size PEG can be prepared. The diols can then be converted to their mono-methoxy derivatives and then activated for conjugation to protein sugar, lipid, nucleotide sugars or DNA/RNA.

Example 1:

Example 1: continuedExample 2:

Example 3:

By varying the ratio of reactants, the base used, temperature, solvent and concentration, one can adjust the reaction to give the predominant size (n) desired.

This approach describes a simple, fast, efficient way to prepare the polyethylene glycols, of any size, in a mono-dispersed size. Purification is simplified by this approach because the difference in size and therefore each molecule's physico-chemical characteristics is very different. This allows the use of simple, standard purification techniques such as silica gel, reversed phase, cellulose, membrane filtration (nano-filtration and ultra-filtration) to be used. The purified PEG diols can then be derivatized into any functional form that is desired.

While this invention has been disclosed with reference to specific embodiments, it is apparent that other embodiments and variations of this invention may be devised by others skilled in the art without departing from the true spirit and scope of the invention.

This Page is inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ BLACK BORDERS

☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLORED OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REPERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images
problems checked, please do not report the
problems to the IFW Image Problem Mailbox**